










Syncope


1. Assess for mechanism of injury, if trauma sustained, refer to **General Trauma Protocol**.
2. Follow **General Pre-hospital Care Protocol**.
3. Position patient
 - A. If third trimester pregnancy, position patient left lateral recumbent.
 - B. Supine for all other patients
4. If patient's mental status remains altered, refer to **Altered Mental Status Protocol**.
-  5. For signs of dehydration or hypotension, administer NS IV fluid bolus.
 - A. Adults up to 1 liter
 -  B. Pediatrics up to 20 mL/kg
 - C. Titrate to normotensive BP
-  6. Obtain 12-lead ECG per **12 Lead ECG Procedure** (May be a basic skill based on MCA selection). If ECG indicates cardiac event or dysrhythmia, refer to Appropriate Cardiac Protocol.
-   7. Additional IV fluids as ordered.



Follow General Prehospital Care Protocol



 For Signs of Hypotension, administer NS IV fluid bolus

- Adults up to 1 liter
-  • Pediatrics up to 20 ml/kg
- Titrate to Normotensive BP

 Obtain 12 Lead ECG per 12 Lead ECG Procedure
 If cardiac event or dysrhythmia, refer to Appropriate Cardiac Protocol

  Additional IV Fluids as Ordered