

Hepatitis Panel/Acute Hepatitis Panel (80074) – NCD 190.33

Indications:
1. To detect viral hepatitis infection when there are abnormal liver function test results, with or without signs or symptoms of hepatitis.
2. Prior to and after liver transplantation.

Limitations:
After a hepatitis diagnosis is established, only individual tests are needed.

Most Common Diagnoses (which meet medical necessity) *	
B15.0 through B17.9	All acute viral hepatitis types
B18.0 through B19.19	All chronic viral hepatitis types
K72.90	Hepatic failure
K74.60	Cirrhosis of liver
K75.81	Nonalcoholic steatohepatitis (NASH)
K75.9	Inflammatory liver disease
R10.0	Acute abdomen
R10.11	Right upper quadrant abdominal pain
R10.12	Left upper quadrant abdominal pain
R10.13	Epigastric pain
R10.9	Abdominal pain
R11.0	Nausea
R11.10	Vomiting
R11.2	Nausea with vomiting
R16.0	Hepatomegaly
R17	Jaundice
R53.1	Weakness
R53.81	Malaise
R53.82	Chronic fatigue
R53.83	Fatigue
R74.01	Elevation of levels of liver transaminase levels
R94.5	Abnormal results of liver function studies

*For the full list of diagnoses that meet medical necessity see the Hepatitis Panel National Coverage Determination 190.33 document.

The above CMS and WPS-GHA guidelines are current as of: 04/01/2025.