



PORT HURON

IMAGE GUIDED PROCEDURES
MEDICAL IMAGING ORDER FORM

TO SCHEDULE A PROCEDURE PLEASE CALL:

Patient Access

(810) 989-1066
Fax: (810) 987-6342

QUESTIONS/CONCERNS PLEASE CALL:

Radiology Nurses

(810) 989-3180
Fax: (810) 989-3197
Hours: Monday - Friday
8:00 am - 4:30 pm

Patient Name: Allergies:

Birthdate: Date/Time of Procedure:

Table with 2 columns: Reason for Exam and Exam Requested. Includes various medical conditions and procedures with checkboxes.

REQUIRED LABS

- ALL PATIENTS ON COUMADIN - PT/INR - to be done day of procedure
For Prep B or C - PT/INR, PLATELETS - 3 -10 days prior to procedure
For Prep E - PT/INR, PLATELETS, TYPE AND SCREEN 2 UNITS PRBC (KIDNEY/RENAL BIOPSY FOR FUNCTION ONLY) must be done at McLaren Port Huron within 10 days of biopsy

Physician Signature: Date:

Attention Patient: Please see reverse side for exam preparation.

McLaren Port Huron
Patient Prep Instructions for Image Guided Procedures

Prep A:

1. Nothing to eat or drink for 2 hours before procedure time.
2. You will take all your usual medications unless you are taking **Coumadin**. Coumadin must be stopped for 5 days prior to your procedure and all lab work will be drawn the day of your procedure. Confirm with your Doctor that it is safe for you to stop taking your Coumadin for this length of time.

Prep B:

1. Nothing to eat after midnight.
2. You may have clear liquids only up until 2 hours before your procedure time. Examples of clear liquids would be black coffee/tea, water, apple juice/cranberry juice – **no milks or pulpy fruit juices**.
3. You will take all your usual medication, except for:
 - **Diabetic medications** – check with your doctor on the need to adjust diabetic medications for the procedure day only.
 - **Coumadin** – must be stopped for 5 days prior to your procedure and all lab work will be drawn the day of your procedure. Confirm with your doctor that it is safe for you to stop taking your Coumadin for this length of time.
 - **Plavix, Aspirin/ASA, or any product containing Aspirin/ASA** – must be stopped for 7 days before your procedure. Confirm with your Doctor that it is safe for you to be off these medications for that length of time.
4. Your lab work (PT, INR, and platelets) must be done 3-10 days before the procedure, except for those taking Coumadin. If you are on Coumadin all lab work will be done the day of the procedure.

Prep C:

1. Nothing to eat or drink for 2 hours before procedure time.
2. You will take all your usual medication, except for:
 - **Coumadin** – must be stopped for 5 days prior to your procedure and all lab work will be drawn the day of your procedure. Confirm with your doctor that it is safe for you to stop taking your Coumadin for this length of time.
 - **Plavix, Aspirin/ASA, or any product containing Aspirin/ASA** – must be stopped for 7 days before your procedure. Confirm with your Doctor that it is safe for you to be off these medications for that length of time.
3. Your lab work (PT, INR, and platelets) must be done 3-10 days before the procedure, except for those taking Coumadin. If you are on Coumadin all lab work will be done the day of the procedure.

Prep D:

1. Nothing to eat or drink for 2 hours before procedure time.
2. You will take all your usual medication, except for:
 - **Coumadin** – must be stopped for 5 days prior to your procedure and all lab work will be drawn the day of your procedure. Confirm with your doctor that it is safe for you to stop taking your Coumadin for this length of time.
 - **Plavix, Aspirin/ASA, or any product containing Aspirin/ASA** – must be stopped for 7 days before your procedure. Confirm with your Doctor that it is safe for you to be off these medications for that length of time.
3. You do not require any lab work for this procedure, except for those taking Coumadin. If you are on Coumadin all lab work will be done the day of the procedure.
4. Your doctor will give you a prescription for Cipro; take as directed the evening before, morning of and evening after the biopsy. You may also receive an order for an enema, take as directed.

Prep E:

1. Nothing to eat after midnight.
2. You may have clear liquids only up until 2 hours before your procedure time. Examples of clear liquids would be black coffee/tea, water, apple juice/cranberry juice – **no milks or pulpy fruit juices**.
3. You will take all your usual medications, except for:
 - **Coumadin** – must be stopped for 5 days prior to your procedure and all lab work will be drawn the day of your procedure. Confirm with your doctor that it is safe for you to stop taking your Coumadin for this length of time.
 - **Plavix, Aspirin/ASA, or any product containing Aspirin/ASA** – must be stopped for 7 days before your procedure. Confirm with your doctor that it is safe for you to be off these medications for that length of time.
4. Your lab work (PT, INR and platelets, *type, and screen for two units PRBC*) must be done at McLaren Port Huron 3-10 days before your procedure, except for those taking Coumadin. If you are on Coumadin all lab work will be done the day of the procedure.